



**Original Article**

**Effect of Dietary Inclusion of Gum Arabic (*Acacia senegal*) on Performance and Blood Chemistry of Broiler Chicks**

Siham, A. Abdalla<sup>1</sup>, Khadiga A. Abdel- Atti<sup>2</sup>, Huwaida E.E. Malik <sup>1</sup>, Bakheit M. Dousa<sup>1</sup> and Khalid M. Elamin<sup>1,\*</sup>

<sup>1</sup> Department of Poultry Production, Faculty of Animal Production, University of Khartoum, Sudan

<sup>2</sup> Department of Animal Nutrition, Faculty of Animal Production, University of Khartoum, Sudan

**ARTICLE INFO**

**Corresponding Author:**

Khalid M. Elamin  
khalid1130@yahoo.com

**How to Cite this Article:**

Abdalla, S.A., K.A. Abdel-Atti, H.E.E. Malik, B.M. Dousa and K.M. Elamin. 2015. Effect of Dietary Inclusion of Gum Arabic (*Acacia senegal*) on Performance and Blood Chemistry of Broiler Chicks. *Global Journal of Animal Scientific Research*. 3(2):305-310.

**Article History:**

Received: 1 December 2014  
Revised: 17 December 2014  
Accepted: 20 December 2014

**ABSTRACT**

The aim of this study was to investigate the effect of feeding graded levels of gum Arabic (0%, 2.5%, %5 and 7.5%) on the performance and blood chemistry of broiler chicks. A total of 192 unsexed broiler chicks (Cobb strain) were randomly assigned in four dietary treatments. Each treatment contained six replicates (8 birds/replicate). Feed and water were offered *ad-libitum* throughout the experimental Period (42 days).. Feed intake, weight gain and feed conversion ratio were not influenced by the dietary inclusion of gum Arabic. The results also revealed that the different levels of gum Arabic had no effect on serum constituents, except total protein which increased significantly. Cholesterol, creatinine and triglyceride levels were decreased significantly in bird fed 5% and 7.5% gum arabic. It is concluded that the incorporation of gum Arabic in broiler chickens diets has a positive effects on broiler chicks performance and reduced serum cholesterol levels, creatinine and triglyceride.

**Keywords:** serum, Cholesterol, creatinine, triglyceride.

Copyright © 2015, World Science and Research Publishing. All rights reserved.

**INTRODUCTION**

Gum Arabic is the dried exudates obtained from stems and branches of *Acacia Senegalor* *Acacia Seyal*. These plants are cultivated in Sudan as a cash crop in agro-forestry systems (Duke, 1981). In Sudan gum Arabic production covers a wide area as it considered as a principle national product .Gum Arabic has been used in many industries such as processed food, dairy products and pharmaceutical preparation e.g. emulsion and pills (Leung. 1980). It was used in folk medicine to treat different diseases. Gum Arabic contains high molecular weight lipoprotein and low molecular weight heterogeneous gum polysaccharides (Nasir,2008). It is indicated that the supplementation with gum Arabic increased nitrogen concentration in renal failure patients consuming a low protein diet (Bliss *et al.*, 1996). In

birds soluble dietary fiber such as oligofructose and inulin were fermented in the cecum (Cehen *et al.*, 2005). However, there are no limitations to the effect of supplementing gum Arabic in layers diet on serum and yolk albumin level. The use of gum Arabic as a feed additive met the safety demand suggested by the international food safety (Androsen, 1986). The objective of the present study was to determine and to assess the effect of dietary supplementation of gum abaci on broiler growth performance on some blood constitutes and metabolites (protein, glucose, cholesterol, creatinine, albumin, calcium, phosphorus and triglyceride).

## MATERIALS AND METHODS

### Experimental Birds and Management

One hundred and ninety two one day old unsexed commercial strain (Cobb) broiler chicks were randomly assigned to four experimental diets. Forty eight birds were used per treatment in a completely randomized design; each treatment was further subdivided into six replicates (8birds/ pen). The initial weight of the chicks in each pen was approximately equal. The chicks were reared in deep litter with feed and water offered *ad-libitum*. The experimental birds were vaccinated against Newcastle disease at 14 day-old (IB) and at the 28 day (Iasota). Gumboro disease vaccine was given at the 21 day. Vitamins were offered for three days after vaccination as supportive dose in drinking water.

### Experimental Diet:

Four starters and four finisher diets were formulated according to nutrient requirement suggested by NRC (1994).The control diet (A) contained 0% gum Arabic while the other three diets B, C and D contained 2.5, 5 and 7.5% gum Arabic respectively. Chicks were fed starter diets at the first three weeks and afterward they were placed on finisher experimental diets. The proximate analysis of gum Arabic was shown in Table 1. The compositions and proximate analysis of gum Arabic starter and finisher diets were shown in Table 2 and 3.

**Table 1: Chemical composition of gum Arabic**

Component	Percentage
Dry mater	87
Crude protein %	2.5
Ash %	3.6
Crude fiber %	0
Arabinose	25
Lactose	14
Methyl glucogonic acid	1.5
pH	4.4
ME kcal/kg diet	3000

### Blood Chemistry

Serum glucose was determined using GOD – POD method as described by (Trinder, 1969) and total serum protein was determined by Biuret method as described by King and Wooton (1965). Cholesterol concentration was determined according to the methods described by Naito (1984).Serum triglycerides were determined using GPO–POD methods as described by Bucolo *et al.*, (1973) and serum calcium was determined by calorimetric method as described by Trinder (1967).Inorganic phosphorus was obtained by the colorimetric method using a kit (Randox laboratory - London) and creatinine was determined using commercial kits (Biosystem reagents Reagent and Instruments).

**Table 2: Composition of experimental broiler starter diets containing different levels of gum Arabic**

Feed Ingredient %	Levels of gum Arabic			
	0.00	2.50	5.00	7.50
Sorghum	60	56.95	53.56	49.95
Groundnut meal	24.2	17.55	15.91	23
Sesame meal	5.5	14	16.33	10
Wheat bran	1.56	0.4	0.4	0.4
Super concentrate*	5.00	5.00	5.00	5.00
Gum Arabic	0.00	2.5	5.00	7.5
Di-calcium phosphate	0.74	0.7	0.69	0.74
Oyster shell	0.24	0.24	0.24	0.24
Salt	0.20	0.20	0.20	0.20
Premix**	0.25	0.25	0.25	0.25
Lysine	0.12	0.12	0.12	0.06
Methionine	0.09	0.09	0.02	0.06
Vegetable oil	2.1	2	2.28	2.6
<b>Calculated Analysis</b>				
ME kcal/kg diet	3197.02	3190.38	3193.3	3190.71
Crude protein %	23.037	23.13	23.00	23.04
Crude fiber %	4.59	4.42	4.36	4.44
Calcium %	1.04	1.16	1.19	1.12
Total phosphorus %	0.45	0.45	0.45	0.45
Lysine %	1.22	1.20	1.19	1.15
Methionine %	0.51	0.58	0.52	0.51

\*contains (%): CP 32, CF 2, Ca 7, P 5, Lysine 11, Methionine 3.7, and ME 1900 kcal/kg.

\*\*provided per kg of diets: vitamin A 8000 IU, vitamin D3 1400 IU, vitamin E 2IU, vitamin K3 2 mg, vitamin B2 4 mg, vitamin B1 2 mg, Ca – d – pantothenate 5 mg, Nicotinamide 15 mg, Choline chloride 100 mg, Folic acid 0.5 mg, vitamin B12 5mcg, Iron 22 mg, Manganese 33 mg, Copper 2.2 mg, Cobalt 0.5 mg, Zinc 25 mg, Iodine 1.1mg.

**Table 3: Composition of experimental broiler finisher diets containing different levels of gum Arabic**

Feed Ingredient %	Levels of gum Arabic			
	0.00	2.50	5.00	7.50
Sorghum	64.82	64.26	62.42	63.61
Groundnut meal	12	13.6	14	15
Sesame meal	7	7	7	7
Wheat bran	8.19	5	4	0
Super concentrate*	5.00	5.00	5.00	5.00
Gum Arabic	0.00	2.5	5.00	7.5
Di-calcium phosphate	0.35	0.4	0.4	0.45
Oyster shell	0.2	0.16	0.15	0.15
Salt	0.20	0.20	0.20	0.20
Premix**	0.20	0.20	0.20	0.20
Lysine	0	0.08	0.08	0.1
Methionine	0	0	0	0
Vegetable oil	2.04	1.6	1.55	0.79
Anticoccidia	0.002	0.002	0.002	0.002
Antifungal	0.1	0.1	0.1	0.1
<b>Calculated Analysis</b>				
ME kcal/kg diet	3190.243	3190.7	3190.20	3190.9
Crude protein %	20.09	20.24	20.07	20.05
Crude fiber %	4.50	4.24	4.100	3.708
Calcium %	0.91	0.91	0.91	0.92
Total phosphorus %	0.38	0.39	0.38	0.39
Lysine %	1.01	1.09	1.08	1.09
Methionine %	0.41	0.40	0.40	0.40

\*contains (%): CP 32, CF 2, Ca 7, P 5, Lysine 11, Methionine 3.7, and ME 1900 kcal/kg.

\*\*provided per kg of diets: vitamin A 8000 IU, vitamin D3 1400 IU, vitamin E 2IU, vitamin K3 2 mg, vitamin B2 4 mg, vitamin B1 2 mg, Ca – d – pantothenate 5 mg, Nicotinamide 15 mg, Choline chloride 100 mg, Folic acid 0.5 mg, vitamin B12 5mcg, Iron 22 mg, Manganese 33 mg, Copper 2.2 mg, Cobalt 0.5 mg, Zinc 25 mg, Iodine 1.1mg.

## Data Collection

At the end of the experimental period two birds from each replicate of each dietary treatment were randomly selected and weighed individually then slaughtered and allowed to bleed. Samples of blood were collected from two birds/replicate (12 birds/treatment) into clean dry test tubes and allowed to clot and serum was separated, collected and analyzed.

## Experimental Design and Statistical Analysis

The statistical analysis for recorded data was carried out using analysis of variance for completely randomized design by general linear model using Statistic program version 8. Means separation were done by LSD and the values were expressed as means and standard error. The means considered significantly different when ( $P < 0.05$ ).

## RESULTS AND DISCUSSIONS

The performance of broiler chicks fed graded levels of powder gum Arabic (0, 2.5, 5 and 7.5%) are presented in Table 4. The results showed that there were no significant differences between dietary treatments in feed intake, body weight gain and feed conversion ratio. Feed intake recorded the lowest value for birds fed control diet (3269.3 g) and the highest value (3628.2 g for the group received 7.5 % gum Arabic diets. Reported feed intake is lower than the results reported by Ali *et al.*, (2013) and Kur *et al.*, (2013). The best results of weight gain were obtained in birds fed 7.5% gum Arabic (1741.3 g). Results for weight gain were higher than the results reported by Ali *et al.*, (2013) and Kur *et al.*, (2013). Feed conversion ratio recorded high results for birds received control diet and 2.5% gum Arabic (1.93 and 1.96 Vs 2.17 and 2.08 for diet 5 % and 7.5% gum Arabic). The findings of the present study were in agreement with the result reported by El-kheir (2009) who found that bird fed high level of gum Arabic recorded increased feed intake. He suggested that increased gum level may enhance the palatability of the diet.

**Table 4: Effect of dietary gum Arabic levels on overall performance of broiler chicks**

Item	Treatment				±SEM
	Gum Arabic level %				
	0	2.5	5	7.5	
Feed intake (g)	3269.3 <sup>b</sup>	3336.3 <sup>b</sup>	3478.2 <sup>ab</sup>	3628.2 <sup>a</sup>	134.32
Weight gain(g)	1698.3 <sup>a</sup>	1703.9 <sup>a</sup>	1615.3 <sup>a</sup>	1741.3 <sup>a</sup>	89.31
F.C.R	1.93 <sup>b</sup>	1.96 <sup>b</sup>	2.17 <sup>a</sup>	2.08 <sup>ab</sup>	0.091

<sup>a-b</sup> = Means with different superscripts in the same row were significantly different ( $P < 0.05$ ).

SEM = Standard error of treatment means

The effects of gum Arabic on broilers blood chemistry are shown in Table 5. Birds fed 5% and 7.5 % gum Arabic obtained lowest level of serum cholesterol compared to control and 2.5% groups (50.74 and 57.76 Versus 68.15 and 67.08 mg/dl) this result in line with El-kheir (2009) who reported that gum Arabic reduced serum cholesterol may be due to its effect by interference with dietary cholesterol absorption. Moreover, Alothman (1998) reported that gum Arabic was observed to be effective in lowering the total plasma cholesterol level compared with insoluble fiber. In the current study diet contained 5% gum Arabic recorded the lowest levels of serum glucose (129.6mg/dl) compared with other groups 57.76-68.15 mg/dl). This result is in agreement with Wadood *et al.*, (1989) who reported that gum Arabic significantly reduces glucose concentration. In the present study serum albumin (2.28 -2.63 g/L), phosphorus (5.03 -5.18 mg/dl), calcium (8.03 – 8.43 mg/dl) and creatinine (0.83-1.35 mg/ dl) levels were not significantly affected by dietary treatments. However Kawase *et al.*, (2007) revealed that the efficiency of Ca absorption in rats was improved by using gum Arabic. In the present study creatinine level numerically decreased for birds fed 5% and 7.5% gum Arabic. These results were in line with the findings of Suliman (2000) who reported that supplemented of gum Arabic orally to patient with chronic renal failure, and it was claimed that it helps to reduce urea and creatinine plasma concentration. Serum triglyceride was decreased by inclusion of gum Arabic 39.88 – 44.58 Vs 57.73 mg/dl in the control group). This result is in line with results obtained by AbdelWahed (2010), Topping *et al.*, (1985) who observed that supplemented dietary level of gum Arabic resulted significant decrease in serum triglyceride. In present study birds fed 2.5% gum Arabic recorded the lowest result for total

proteins (2.72 g/l) while the other groups were not significantly different for the trait (3.12-3.14 g/l).

**Table 5: Effect of dietary gum Arabic levels on serum constituents of broiler chicks**

Item	Treatment				±SEM
	Gum Arabic level %				
	0	2.5	5	7.5	
Cholesterol(mg/dl)	68.15 <sup>a</sup>	67.08 <sup>a</sup>	50.74 <sup>b</sup>	57.76 <sup>ab</sup>	3.93
Albumin (g/l)	2.37 <sup>a</sup>	2.28 <sup>a</sup>	2.63 <sup>a</sup>	2.38 <sup>a</sup>	0.19
Glucose (mg/dl)	158.8 <sup>a</sup>	138.08 <sup>ab</sup>	129.6 <sup>b</sup>	139 <sup>ab</sup>	8.48
Triglyceride (mg/dl)	57.73 <sup>a</sup>	39.88 <sup>b</sup>	44.58 <sup>ab</sup>	40.38 <sup>b</sup>	5.45
Total protein (g/l)	3.12 <sup>a</sup>	2.72 <sup>b</sup>	3.13 <sup>a</sup>	3.14 <sup>a</sup>	0.21
Phosphorus (mg/dl)	5.18 <sup>a</sup>	5.03 <sup>a</sup>	5.12 <sup>a</sup>	5.03 <sup>a</sup>	0.28
Calcium (mg/dl)	8.6 <sup>a</sup>	8.03 <sup>b</sup>	8.43 <sup>ab</sup>	8.38 <sup>ab</sup>	0.37
Creatinine (mg/dl)	1.35 <sup>a</sup>	1.33 <sup>a</sup>	0.92 <sup>ab</sup>	0.83 <sup>b</sup>	0.32

<sup>a - b</sup> = Means with different superscripts in the same row were significantly different (P 0.05). SEM = Standard error of treatment means. Values are means of 6 replicates per treatment (2 birds/replicate)

## CONCLUSION

It is concluded that the incorporation of gum Arabic in broiler chickens diets has a positive effects on broiler chick performance and on the reduction of serum cholesterol levels, creatinine and triglyceride.

## REFERENCE

- Abdelwahed, N., O.F. Idris and H.I. Sari. 2010. The effect of feeding gum Arabic on serum total lipoprotein cholesterol in hypercholesterolemic rats. *Assiut Veterinary Medicine*. 57: 128.
- Ali, F.M., K.A. Abdel Atti, B.M. Dousa, K.M. Elamin. 2013. Performance and blood chemistry of broiler birds fed different levels of dietary Rashad (*Lepidium sativum*). *Wayamba journal of Animal science*. No. 1368709198. <http://www.wayambajournal.com/paper.php?n=1368709198>.
- Al-Othman, A.A.,R.A. Al-Shargawi, F.M. Hewedy, and M. Hamdi. 1998. Plasma total, lipoprotein cholesterol, organs cholesterol and growth performance in rats fed dietary gum Arabic. *Food Chemistry*. 62: 69-72.
- Anderson, D.M. 1986. Evidence of the safety of gum Arabic (*Acacia Senegal* (L) Willd) as a food additive- a brief review. *Food Addit. Contam.* 3:225-230.
- AOAC (Association of Official Analytical Chemist). 1990. Official Methods of Analysis 12<sup>th</sup> ed. Washington. DC.
- Bliss, D.Z.,T.P. Stein, C.R. Schleifer and R.G.Settle. 1996. Effect of gum Arabic on chronic renal failure. *American Journal of Clinical Nutrition*. 63: 392-398.
- Bucolo, G., and M.David. 1973. Quantitative determination of serum triglycerides. *Clinical Chememistry*. 19(5) 476-482.
- Chen, Y.C., C. Nakthong and T. Chen. 2005. Improvement of laying hen performance by dietary prebiotic chicory oligofructose and inulin. *International Journal of Poultry Science*. 4:103-108.
- Duke, J.A. 1981. Hand book of Legumes of world economic importance. Plenum Press. New York.
- El-kheir, M.K.S., K.E.A. Ishag, A.A. Yagoub, and A.A. Abu baker. 2009. Supplementation Laying Hen Diet with Gum Arabic (*Acacia Senegal*): Effect on Egg Production, Shell Thickness and Yolk Content of Cholesterol, Calcium, and Phosphorus. *Asian Journa of poultry Science*. 3(1):9-14.
- Kawase, A.,N. Hirata, M. Tokunage, H.Matsuda. AndM. Iwaki. 2007. Gum Arabic enhances intestinal calcium absorption in rats.*Journal of health Science*. 53: 622-624.
- King, E.S., andJ.G.P. Wooton. 1965. Determination of total protein in plasma or serum. In *Medical Biochemistry*, pp, 138-140. Churchill. London.
- Kur, A.T.Y., K.A. Abdel Atti, B.M. Dousa, H.A.A. Elagib, H.E.E. Malik, K.M. Elamin. 2013. Effect of treated cowpea seeds on broiler chicken. *Global Journal of Animal scientific Research*. 1(1):53-60.
- Leung, A.Y. 1981. Encyclopedia of common natural ingredients used in food, drugs and cosmetics. John Wiley and Sons. New York.
- Naito, H.K. 1984. Determination of plasma and serum cholesterol. *Clinical Chemistry*. pp:1194-1206.

- Nasir, O., F. Artune, A. Saeed, M.A. Kambal, H. Kalbacher, D. Sandulache, K.M. Bioinic, N. Johove and F. Long. 2004. Effect of Gum Arabic (*Acacia senegal*) on water and electrolyte balance in healthy Mice. *Journal of Renal Nutrition*. 18:230-238.
- NRC. 1984. Nutrient Requirement of poultry (seventh edition) National Research Council. Nutritional Academy of Science. Washington DC. USA.
- Suliman, S.M., M.I. Hamdouk, and M.B. Elfaki. 2000. Gum Arabic fiber as a supplement to low protein diet in chronic renal failure patient in Sudan. *Physicians' association 17<sup>th</sup> conference Friend ship hall*, 21-23 March, Khartoum, Sudan.
- Topping, D., R.J. Illman, and R.P. Trimble. 1985. Volatile fatty acid concentration in rats fed diet containing gum Arabic and cellulose separately and a mixture. *Nutrition Report. International*. 32(4): 809-814.
- Trinder, P. 1967. Calorimetric micro-determination of serum calcium. In *Microanalysis in Medical Biochemistry*. (Edited by Wooton, J.D.P.). pp: 76-77. Churchill. Ltd. London.
- Trinder, P. 1969. Plasma and serum glucose determination. *Annual of Clinical Biochemistry*. p: 624.
- Wadood .A.A., N. Wadood and A.S. Shah. 1989. Effect of acacia Arabica and caralluma edulis on blood glucose levels of normal and allaxon diabetic rabbits. *Journal of Pakistan Medical Association*. 39:208-212.