



Review Article

A Review on Dromedary Camel Milk Products and Their Uses

Aleme Asresie^{1,*} and Mulugojjam Adugna²

¹College of Agriculture and Environmental Sciences, Department of Animal Sciences, Adigrat University, Adigrat, Ethiopia

²College of Natural and Computational Sciences, Department of Biology, Debretabor University, Debretabor, Ethiopia

ARTICLE INFO

Corresponding Author:
Aleme Asresie
almasres06@gmail.com

How to cite this article:
Asresie, A., and M. Adugna.
2014. A Review on Dromedary
Camel Milk Products and
Their Uses. *Global Journal of
Animal Scientific Research*.
2(3): 285-290.

Article History:
Received: 20 June 2014
Revised: 1 July 2014
Accepted: 2 July 2014

ABSTRACT

This paper reviewed researches on dromedary camel milk products and their uses. Milk from the lactating camel must provide nourishment for her young calf as well as for human, not a great deal will be left for milk products. Moreover, the composition of camel milk does not allow for making some of the accepted products that are made from cow, sheep and goat milk. Nevertheless, milk products are made from camel milk, and the milk itself is used for purposes other than simply nutrition. Camel products have various names in various parts of the world. In the Caucasus it is called kefir; in Armenia, matzoon; in India, dahdi; in Sardinia, gioddu; in Bulgaria, yoghurt; in Ethiopia, Dhanaan and in Syria, Israel and Egypt, lehben. Some authors describe butter being made from camel milk. While others categorically state that butter cannot be made from camel milk. The preparation of butter from camel milk is not as easy as from milk of other animals owing to its unique milk-fat properties. Soft cheese can be made from camel milk. Camel milk and their product are used therapeutically against dropsy, Jaundice, problems of the spleen, tuberculosis, asthma, anaemia, and piles. In Ethiopia camel milk is considered as having aphrodisiac powers. In Somalia, among the pastoral tribes, it is believed that milk drunk on the night when the camels first drink water, following a long period of thirst, has magical powers. We suggest that consumption of camel milk and their products may have remedial effects for many different life-style related diseases.

Keywords: Aphrodisiac power, dhanaan, dromedary, remedial effect.

Copyright © 2014, World Science and Research Publishing. All rights reserved.

INTRODUCTION

Milk from the lactating camel must provide nourishment for her young calf as well as for human, not a great deal will be left for milk products. Moreover, the composition of camel milk does not allow for making some of the accepted products that are made from cow, sheep and goat milk. Nevertheless, milk products are made from camel milk, and the milk itself is used for purposes other than simply nutrition. In most pastoral communities, where milk for human consumption is obtained from more than one domesticated species of animal, milk products are made after milk of various animals in mixed. It is often unclear therefore if some

of the products can be made from camel milk alone (Dhal and Hjort, 1976). This is often the case when camel milk is mixed with fresh or churned goat milk (Gast *et al.*, 1969). This mixture is made with certain quantities of camel milk added until the required taste is obtained.

Not all communities use camel milk for making products. The milk of the Afar camels in Ethiopia, for instance, is not allowed to be processed (Dhal and Hjort, 1979). On the other hand the milk of the Somali camels in Ethiopia is allowed to be processed into different dairy products for instance butter made from camel milk alone and blending it with goat milk (Asresie *et al.*, 2013). When camel milk is not consumed fresh it must be processed as soon as possible both because its keeping quality seems to be poor and as it is further adversely affected by the climate it soon goes bad if not treated. Many superstitions and beliefs have evolved around camel milk and milking only specific members of the family can graze the animals and the milk is considered as having medicinal as well as mystical properties (Hartley, 1979). Therefore, this manuscript Endeavour's to present a detailed discussion on dromedary camel milk products and their uses.

CAMEL MILK PRODUCTS

Fermented milk products

Under warm conditions raw milk does not keep for long and actually its fermentation appears to be a means to preserve it not only for a limited period of time. Fermented products have various names in various parts of the world (Aggarwala and Sharma, 1961). In the Caucasus it is called kefir; in Armenia, matzoon; in India, dahdi; in Sardinia, gioddu; in Bulgaria, yoghurt; in Ethiopia, Dhanaan and in Syria, Israel and Egypt, lebben. The method of preparation of fermented milk consists in heating the milk to the boiling point (Aggarwala and Sharma, 1961) so as to kill bacteria.

The milk is well stirred and kept overnight at ambient temperature. By next morning it has curdled. At that stage it has acquired a sour taste and the typical flavour of fermented milk has developed. Pathogenic bacteria were killed when the milk was boiled and conditions have developed which will make it difficult for them to develop, assuring that a re-infection took place at a later stage. Therefore, fermented milk products are edible for some time. The method for commercial manufacture of kefir from camels' milk: milk is flashing pasteurized at 85⁰C, getting rid by this means of pathogenic bacteria. The milk is then cooled to 26–30⁰C and then inoculated with a 3 to 6 percent of kefir culture. It is then bottled. After incubation at 20–26⁰C for 8 to 12 hours, a soft coagulum is formed and its acidity reaches 60–70⁰T. The product is then allowed to ripen for 24–28 hours at 6 to 8⁰C. The end product has a refreshing flavour and a thick creamy consistency. It is white and without gas. The acidity of one-day old kefir is in the region of 95⁰T and its alcohol content is about 4 percent (Kheraskov, 1964).

“Chal” or “shubat”, is a white sparkling beverage that has a sour flavour (Lakosa and Shokir, 1964). The “Chal” is prepared by first souring it in a skin bag or ceramic jar, normally with a capacity of 30kg. Previously soured milk is added to the fresh milk. It is well mixed and each day, for 3 to 4 days, fresh milk is added to the mixture. Eventually the end product must have 3 to 5 times the original volume of “Chal” that was initially added. This is the best ratio for the “Chal”. It was found that camel milk does not sour at temperatures below 10⁰C and this for up to 72 hours. At 30⁰C the milk sours in approximately 8 hours, compared with cow's milk, which sours within 3 hours at a temperature of 30⁰C. The comparison between the composition of camel milk and camel “Chal” is as follows (Grigor'yants, 1954): The “Chal” contains Lactobacilli lactic; streptococci and yeast (Kieselev, 1956). “Chal” was successfully prepared by using cultures of Lactobacillus casei, Streptococcus thermophilus and lactose fermenting yeasts and incubating inoculated milk for 8 hours at 25⁰C and subsequently for 16 hours at 20⁰C. Holder pasteurization did not affect the quality of the milk, but pasteurization at 85⁰C for 5 minutes caused the milk to have a bad flavour. “Chal” made

from pure cultures of *Lactobacillus casei*, *Streptococcus thermophilus* and species of *Torula* had markedly less no-fat solids and lactose than the milk from which it was made (Kuliev, 1959).

In Mongolia “Tarag” is a cultured milk product similar to yoghurt, while “Unda” is a product produced by lactic and alcohol fermentation of camel and other animals' milk (Accolas *et al.*, 1975). In the Ahaggar region of the Sahara milk is fermented in a manner similar to that mentioned above (Gast *et al.*, 1969). The animals are milked into special jars, made of Tamari wood, which can hold 2 to 3 litres of milk. The milk is mixed in the jars, and if not immediately drunk, it is stored in containers made of animal skin. In these skin containers the milk sours. The milk of the previous day is mixed with fresh milk until sour. There are great changes in the fat percentages of the milk products. As it is a practice to water the animals only once every 4 to 5 days, and as the lack of drinking water can cause a decrease in fat percentage (Yagil and Etzion, 1980) the fluctuating fat content is understandable. A lebben is also made by sweetening fresh milk (Gast *et al.*, 1969). If water is added to this mixture a longer storage time is achieved. This lebben can be kept for 5–6 days in the summer, and up to 10 days in the winter.

Khoa and other non-fermented whole milk products

“Khoa” is made by evaporating small amounts of milk over a hot, steady fire. (Aggarwala and Sharma, 1961). The milk is continuously stirred to prevent scorching. At first the mass left over has a buttery consistency, but after cooling, it turns into semi-solid dough with a sweet taste. “Khoa” can be kept for about 200 days. If sugar is added it keeps for longer periods. “Rabbri” is also made by heating milk in a shallow iron pan over a hot fire. The difference with “Khoa” is that the solids are removed successively from the thin layer of coagulated milk on the surface. Then the product is allowed to cool. When the milk reaches a fifth to an eighth of the original volume, it is removed from the fire. The mass is now gently mixed, without damaging the flakes that have formed. Sugar is added and it is then allowed to cool.

Butter and derived products

Some authors describe butter being made from camel milk (Asresie *et al.*, 2013; Shalash, 1979) while others categorically state that butter cannot be made from camel milk (Dickson, 1951). The preparation of butter from camel milk is not as easy as from milk of other animals owing to its unique milk-fat properties. The fat is distributed as small micelle-like globules in the milk (Dong-Wei, 1981); Yagil and Etzion, 1980). In addition, the fat is apparently bound to protein and there is a great difference in fatty acid composition. (Gast *et al.*, 1969). Samples of camel butter are characteristically rich in polyunsaturated fatty acids. There are only traces of fatty acids with chains shorter than C-12 lauric acid. The butter does contain normal amounts of C-16 palmitic acid, and has very high content of the polyunsaturated C-18 oleic and linoleic acids, when compared with butter obtained from milk of other animals (Gast *et al.*, 1969). Knoess observed that butter can be obtained after 15 to 20 minutes churning, whereas according to (Shalash, 1979) it can take up to about four hours. Butter can be made by churning fresh camel milk at 24–25⁰C in a special blender (Lakosa and Rakin, 1964). At lower temperatures the cream of the camel milk will not churn. Water content of butter was found to be about 13 percent and its acidity 3.3⁰T. Specific gravity was 0.923 and it melted at 44.5⁰C. This butter is used for cooking and is not eaten as butter. It is sometimes used by women as a cosmetic (Gast *et al.*, 1969).

In the Sahara, butter is made by placing camel milk into a thin, hairless, goat-skin for 12 hours. This skin is never washed with water. Inside the temperature rises to 28⁰C, a temperature very similar to that used in the blender (Lakosa and Rakin, 1964). In winter the goat-skin is often placed into the ground near a warm fire to obtain the optimum temperature before making butter. This aids in the fermentation. Churning is done when the container is

half filled with sour milk. Air is blown into the container and the top is tied off. It is hung on a tent pole and rapidly swung to and fro. This is done in the early morning and the amount of butter obtained is determined by the skill of the man doing the churning. No churning is done during the day as solar heat apparently impedes proper separation. Some cold water is added into the goat-skin before the end of churning. This aids in solidifying the butter. It is then placed in a wooden bowl or kettle.

Fresh butter is not eaten, but is often used as a base for medicines. The fresh butter is difficult to preserve. It is not limpid and becomes rancid rapidly. Most of the butter is thus melted down to make Shmen or "Semma". The butter is melted at 100–120⁰C for 30 minutes. A clarifying agent is added to hot butter and it is stirred with a wooden spoon.

This agent can be crushed dates or a grated, roasted piece of ram horn, or leaves of certain plants or seeds. Heating destroys the bacteria and the clarification agent collects the dirt and floats to the top, where it can be skimmed off. If dates are used as clarifying agent it is then given to children. If not, it is thrown away. The leaves give the specific colouring and aroma to the butter. The Bedouin in the Sinai Peninsula place the camel milk that is left over in a big clay jar, were it allowed to partially ferment. Then they place the milk into a leather container and shake it for about 4 hours and subsequently extract butter. Milk from quite a few camels is normally required to obtain enough butter for the requirements of the herder families. Camel butter has a harder consistency than the butter of sheep milk. The butter has a greasy appearance and taste, so only little is eaten and it is used mainly for cooking.

Cheeses

Soft cheese can be made from camel milk. The fat is bound to the milk proteins and the casein is also different from that of other animals (Ottogalli and Resmini, 1976). The alpha and beta caseins were found to react much more slowly on electrophoresis. Nevertheless, several plants that make rennet coagulated cheese from camel milk exist (Dilanyan, 1959; Mares, 1954). The Tourag nomads on the other hand say that cheeses cannot be made from camel milk, as it does not curdle, and so discussion on cheese making in the Sahara is restricted to cheeses made from milk of other animals (Gast *et al.*, 1969). However, cheeses can also be made by mixing camel milk with milk of other animals (Kheraskov, 1965.; Rao *et al.*, 1970). The cheeses made are often spiced cheeses or made without salt or sugar. The difficulty in making cheeses in the Sahara most probably refers to the technique which is being used. This is apparent as the addition of the peptic enzyme that is collected from rabbit stomach, or from the abomasums of young goats, to camel milk causes the formation of a coagulum. This coagulated mass is soft like cotton wool (Gast *et al.*, 1969).

As previously mentioned, in summer the lack of proper amenities in many countries to keep milk at low temperature does not allow for storing milk. The left-over camel milk is thus curdled and soured. Casein can be prepared from this milk and the product is known as "industrial casein" (Pant and Chandra, 1980), because the product is considered unfit for human consumption. It is used for making glue and gums. Whereas industrial casein and its fractions made from cow milk were found to be rich in crude proteins, the industrial casein and its fractions when made from camel milk are poor in crude proteins. Simple and effective methods were standardized for obtaining proteins from milk whey. The proteins of camel milk whey have a relatively higher percentage of nitrogen than those obtained from the whey of cow milk (Pant and Chandra, 1980). The high percentage of proteins and the amino acid composition of camel milk industrial casein suggest that it could make a satisfactory dietary supplement to the human diet. The unpleasant odour and taste however, make it unsuitable for consumption by man or animals. It is thought that it would be possible to purify the camel industrial casein and so make it fit for human consumption. The Bedouin of the Sinai Peninsula and the Tourag nomads (Gast *et al.*, 1969) make a dry cheese called "Afig". These are balls of cheese that are made from butter milk, after the butter has been made, as mentioned previously. The Afig cheese balls are placed on the sides of the tents to dry out. No

other kinds of cheeses are made. The reason for this is said to be that the camel spoke to the Prophet Mohammed and it was agreed that no cheeses would be made from her milk and that her wool would not be dyed.

MISCELLANEOUS USES OF CAMEL MILK

Medicinal properties

In India camel milk is used therapeutically against dropsy, Jaundice, problems of the spleen, tuberculosis, asthma, anaemia, and piles (Rao *et al.*, 1970). The “Chal” and other lung ailments (Gast *et al.*, 1969.) have proven beneficial in the treatment of tuberculosis (Akundov *et al.*, 1972). A clinic has been established in which milk is used for treatment (Urazakov and Bainazarov, 1974). Patients with chronic hepatitis had improved liver function after being treated with camel milk (Sharmanov *et al.*, 1978). In fact, camel milk was as effective as ass milk and superior to treatment with only medication or a diet consisting of cow milk proteins. The camel milk works as a laxative on people unaccustomed to drinking this milk (Rao *et al.*, 1970). Apparently stomach upsets only occur when the milk is drunk while still warm. When it is cool, no ill effects have been noted (Gast *et al.*, 1969). The milk also apparently has slimming properties (Yasin and Wahid, 1957).

Camel milk is given to the sick, the elderly and the very young because of the belief that it is not only healthier, but works especially well in bone formation (Gast *et al.*, 1969). The belief among the Bedouin of the Sinai Peninsula is that any internal disease can be cured by drinking camel milk. The milk is said to be of such strength, and to have such health-giving properties, that all the bacteria are driven from the body. This said to be true only for camels that eat certain shrubs and bushes. The shrubs and bushes are, themselves, used in the preparation of medicines. However, camels which eat straw are said to lose this ability.

Mystical properties

In Ethiopia camel milk is considered as having aphrodisiac powers (Rao *et al.*, 1970). In Somalia, among the pastoral tribes, it is believed that milk drunk on the night when the camels first drink water, following a long period of thirst, has magical powers (Mares, 1954). “He who drinks milk on this night from a thirst-quenched camel will lose the thorns that have penetrated his feet, even from childhood”. In the Sahara there is a superstition that when camel milk is given to a certain guest, only the milk of one particular camel is given to him. (Gast *et al.*, 1969). Therefore, if the guest casts an evil eye on the herd, only the camel, whose milk he has drunk, will be affected and will stop lactating.

CONCLUSIONS

The consumption of camel milk and their product not considered as a cultural value in the world except in pastoral community as well as in Muslim religious follower country people compared to the consumption of milk and derived products from cow, goat, buffalo and Sheep. As for the camels, it seems that this requirements less fulfilled than for other animals with some negative hypothesis remains, which must be overcome gradually. The authors of this article are engaged to positively review and to further propagandize nutritional and therapeutic value of camel milk and their products. We suggest that camel milk and their products may have remedial effects for many different life-style related diseases, including, hypertension and cardiovascular diseases.

ACKNOWLEDGMENT

I am deeply grateful and indebted to all sources of materials used for reviewed this manuscript have been duly acknowledged.

REFERENCE

- Accolas, J.P., J.P. Deffontaines, F. Aubin, F. Grosclaude, S. Kuzdzal-Savoie, D. Langlois, and R. Jeunet. 1975. Rural activities in the Mongolian Peoples' Republic II Dairy Products (Fr.). *Etudes Mongoles*. 6: 7-98.
- Asresie, A., S. Eyassu, and M. Mohammed. 2013. Churning efficiency and microbial quality of butter made from camel milk alone and blending it with goat milk. *Net. J Agri. Sci. 1(3): 75-80*.
- Aggarwalda, A.C. and R.M. Sharma. 1961. A laboratory manual of milk inspection. Asia Publishing House.
- Akhundov, A.A., B. Dyrdyev, and E.R. Serebryakov. 1972. Effect of combined treatment on water electrolyte exchange in pulmonary TBC patients. *Zdravookhr. Turkm.* 16: 40-44.
- Dahl, G. and A. Hjort. 1976. Having herds: Pastoral herd growth and household economy. Univ. Stockholm.
- Dahl, G. and A. Hjort. 1979. Dromedary pastoralism in Africa and Arabia. In: *Camels IFS Symposium Sudan*. 447- 458.
- Dickson, H.R.P. 1951. *The Arab of the Desert*. London, George Allen & Unwin Ltd. 409-446.
- Dilanyan, Z. and I. Gabrielyan. 1959. The Significance of Some Physico-Chemical Properties of Milk in Cheese Making. *Molochnaya. Prom., U.S.S.R* 10: 33.
- Dong Wei. 1981. Chinese camels and their productivities. In: *IFS Symposium Camels. Sudan*. 55-72, Chinese camels' Personal communication.
- Gast, M., L. Mauboisj, and J. Adda. 1969. Le lait et les produits laitiers en Ahaggar. *Centr. Rech. Anthr. Prehist. Ethn.*
- Grigoryants, N.N. 1954. Composition of camel milk and chal. *Vop. Pit. in Russian*. 13: 41-5.
- Hartley, J.B. 1979. Camels in the Horn of Africa. In: *Camels. IFS Symposium, Sudan*, 109-124.
- Kheraskov, S.G. 1961. Composition, properties and nutritive value of camels' milk. *Vop. Pitan.* 20: 69-72.
- Kheraskov, S.G. 1965. Camel milk - a valuable food product. *Konevod. Konny Sport.* 35:14-15.
- Kieselev, N. 1956. Bacteriological examination of chal (Ru). *Mol. Prom.* 17: 31-34.
- Kuliev, K. 1959. The utilization of camels' milk. *Mol. Promyslenn* 20: 28.
- Lakosa, I.I. and A. Shokin. 1964. Milk production. In: *Camel Science Technical Agricultural Publication*. Kolov, Moscow. 113-120.
- Mares, R.G. 1954. Animal husbandry, animal industry and animal disease in the Somaliland Protectorate. *Brit. Vet. J.* 110: 411-423.
- Ottogalli, G. and P. Resmini, 1976. Notizie sui problemi lattiero-caseari della Somalia. *Alcuni datusulle caratteristiche chimico-fisiche del latte di zebu e di dromedaria l'industria del latte*. 12: 3-10.
- Pant, R. and P. Chandra. 1981. Composition of cow and camel milk proteins and industrial casein. *Dairy Sci. Abstr.* 43(3):1763
- Rao, M.B., R.C. Gupta, and N.N. Dastur. 1970. Camels' milk and milk products. *Ind. J. Dairy Sci.* 23: 71-78.
- Shalash, M.R. 1979. Utilisation of camel meat and milk in human nourishment. In: *IFD Symposium. Camels Sudan*. 285-306.
- Sharmanov, T.Sh., R.Kh. Kadyrova, O.E. Shlygina, and R.D. Zhaksylykova. 1978. Changes in the indicators of radioactive isotope studies of the liver of patients with chronic hepatitis during treatment with whole camels' and mares' milk. *Voprosy Pitaniya 1: 9-13*.
- Urazakov, N.U. and S.H. Bainazarov. 1974. The 1st clinic in history for the treatment of pulmonary tuberculosis with camel's sour milk. *Probl. Tuberk.* 2: 89-90.
- Yagil, R. and Z. Etzion. 1980. The effect of drought conditions on the quality of camels' milk. *J. Dairy Res.* 47: 159-166.
- Yasin, S.A. and A. Wahid. 1957. Pakistan camels. A preliminary survey. *Agric. Pakist.* 8: 289-297.